

NORTH MISSISSIPPI HEALTH SERVICES

COVID-19 Treatment • Ambulatory Referral

Patient's name	SSN#	Date of birth
CityState	Contact phone number	
Order: COVID-19 antibody infusion per protocol. Drug choi availability. May be given IV or SubQ.	ice will be based on current EUA	, NMMC committee recommendations and
Observe for one (1) hour. Initiate NMMC Outpatien		for emergency.
Pharmacist to determine solution and rate based on Vital Signs including O ₂ Sat: Obtain at start of infusion, repeat in		eat after 1 hour of post infusion monitoring
For SubQ administration, vital signs will be done upon admission		cat and 1 nour of post-infusion momoning.
Patient Inclusion Criteria		
Date of symptoms onset Date of Outpatient COVID POSITIVE Test (mild to	moderate symptoms by the NIH	(criteria)
Symptom onset within the last 10 days	moderate symptoms by the twin	(cincina)
Inclusion criteria:		
Outpatient COVID positive patient (mild to moderate	te symptoms by the NIH criteria)	
Symptom onset within last 10 days		
$\Box \operatorname{Age} \ge 18$		
 Are high-risk for progressing to severe COVID-19 a Patient must have oxygen saturation greater than or 	-	
If patient is on O_2 , they must not have an increased oxygen requirements OR		
If the patient has a known baseline pulse ox of less than 94% and does not meet the criteria for home oxygen, the provider must		
document that in his/her note for the patient to be eligible.		
High risk is defined as patients who meet at least one of	the following criteria:	
$\Box \text{ Aged } \ge 65 \text{ years}$		
 Obesity (BMI >35) Chronic lung diseases (e.g., chronic obstructive pulmonary disease, moderate-to-severe asthma, interstitial lung disease, 		
cystic fibrosis, pulmonary hypertension)	monary disease, moderate-to-sev	ere asunna, interstitiar fung disease,
An immunocompromising condition (e.g. organ or b steroids) or immunosuppressive treatment	oone marrow transplant, severe in	nmunocompromising therapy, excluding
Chronic liver diseases (e.g., cirrhosis, NAFLD, alco	holic liver disease, autoimmune	hepatitis)
CKD stage $4/5$ (GFR < 30) and ESRD		
 Active Tuberculosis Pregnancy 		
 I regnancy Neurodevelopmental disorders (e.g., cerebral palsy) 	or other conditions that confer n	nedical complexity (e.g., genetic or metabolic
syndromes and severe congenital anomalies)		leanen comprenny (eig., genere er mensene
Medical-related technological dependence (e.g., trac COVID-19)	cheostomy, gastrostomy, or positi	ive pressure ventilation that is not related to
\square Aged >40 years, and at least one of the following:		
Diabetes		disorder (e.g., mood disorders, anxiety/fear
Cardio- (and cerebro-)vascular disease	spectrum disc	ers, substance use disorder, schizophrenia
Sickle cell disease	Obesity (BMI	
Smoking		
See CDC website https://www.cdc.gov/coronavirus/201	9-ncov/need-extra-precautions/p	eople-with-medical-conditions.html
Exclusion Criteria:		
• Do NOT meet inclusion criteria		
 Weight < 88 lbs Require oxygen due to COVID-19 or who require an indication of the second second	crease in oxygen from baseline	
 History of allergic reaction to monoclonal antibody adm 		
Fact Sheet for Patients, Parents, and Caregivers given to patient.		
YES NO		
I acknowledge that I have read and understand the training		
https://www.nmhs.net/medical-professionals/covid-infusion	n-treatment/	